



Targeting protein kinases in cancer therapy: a success?

Mark A Pearson[†] and Dorian Fabbro

The fundamental role of kinases in cancer progression has promoted the development of a plethora of therapeutic inhibitors. Despite the promise of effective treatment with little associated toxicity, the clinical experience with these agents has been mixed. This review will summarize recent advances made in the development of kinase inhibitors to highlight emerging issues and the strategies by which they are being addressed.

Expert Rev. Anticancer Ther. 4(6), 1113–1124 (2004)

CONTENTS

Rationale for targeting kinases in cancer therapy

Biology & target validation

Selectivity & approaches

Clinical examples of kinase inhibitors

Summary & conclusions

Expert opinion

Five-year view

Key issues

References

Affiliations

Rationale for targeting kinases in cancer therapy

Many of the defining characteristics of cancer, including uncontrolled growth, survival, neovascularization, metastasis and invasion, result from perturbation of regulatory signaling pathways, which are normally under tight control. As advances are made in understanding the mechanisms underlying the development of cancer, it has become clear that particular pathways are more frequently deregulated. Deregulation, whether as a result of deletion, mutation or amplification of component gene products, is manifested as aberrant activation of key regulators of these pathways, a prime example of which are kinases [1]. A molecular targeted therapy that specifically inhibits such pivotal regulators would be expected to bypass the toxicity associated with currently used chemotherapeutic agents. This possibility has spurred intense activity within the pharmaceutical industry into the development of agents directed against protein kinases. Currently, over 20 different kinases, the majority being receptor tyrosine kinases (RTKs), are being considered as potential therapeutic targets in oncology.

Although the success of imatinib and trastuzumab has provided a proof of concept that such agents can be both therapeutically effective and retain an acceptable safety profile, the clinical experience with many tyrosine kinase inhibitors has been less promising. This review is not intended as a general overview of the role of kinases in cancer or kinase inhibitors

currently in the clinic, but instead will focus on the emerging criteria that should be met for a kinase inhibitor to be successful.

Biology & target validation

Understanding the role of a potential target in cancer development and progression is as relevant as the efficient optimization of an inhibitory compound's potency, toxicity and pharmacokinetic profile. To be a valid target, a kinase should play a fundamental role in the pathogenesis of the disease. Activating mutations have been used as a rationale to determine potential kinase and disease targets in oncology. Genetic screening of families has linked germline kinase mutations, which lead to a gain of function, with a predisposition to cancer, for example, the association of mutations in c-Met with familial papillary renal cell carcinoma or multiple endocrine neoplasia Type 2 in families with c-Ret mutations [2], or gastrointestinal stromal tumor (GIST) with activating c-Kit mutations [3]. The presence of identical mutations in sporadic versions of the same cancer further reinforces the hypothesis of a fundamental role in progression of the disease. Increased understanding of the molecular history of the disease is also invaluable. Mutations that are found to be expressed in early phases of the disease, such as those found within c-Kit in GIST or the Philadelphia chromosome translocation in chronic phase chronic myeloid leukemia (CML), are more likely to drive disease progression than those that occur later. A correlation between

[†] Author for correspondence
Novartis Institutes for Biomedical
Research Basel, Novartis Pharma AG,
WKL 125.4.16, CH-4002,
Basel, Switzerland
Tel.: +41 616 961 016
Fax: +41 616 963 835
mark_andrew.pearson@pharma.
novartis.com

KEYWORDS:
antibody, biomarker,
clinical trial, kinase inhibitor,
patient selection, resistance

the metastasis/aggressiveness of a type of cancer and a particular mutation has also been used to validate a candidate kinase, for example, epidermal growth factor receptor (EGFR)vIII mutants in glioblastoma [4]. Accumulating evidence suggests that the prevalence of kinase mutations in solid tumors is far greater than previously thought. A large-scale genomics sequencing approach identified an unexpectedly high frequency of BRAF mutations in melanoma [5]. In addition, point mutations in several tyrosine kinases and within the phosphatidylinositol 3-kinase catalytic subunit (PI3KCA) have been identified in colorectal cancer. Mutations in PI3KCA, that resulted in higher kinase activity, were also found to occur within several other cancer types, including glioblastoma and gastric cancer [6,7].

In the absence of information on kinase mutational status, overexpression of a kinase, particularly RTKs, has been used as a rationale to design inhibitors, for example, EGFR-targeting agents such as gefitinib and erlotinib. However, it is worth noting that a low ($\leq 10\%$) percentage of patients responded to treatment with these agents and this response has been linked to the presence of novel mutations in the EGFR [8,9]. Despite this, it is clear that kinase inhibitors can be clinically effective in situations where the kinase is not mutated (e.g., treatment of patients with breast cancer overexpressing HER-2 with trastuzumab or dermatofibrosarcoma protuberans with imatinib) [10,11]. One of the current priorities in kinase-centric cancer research is therefore the identification of additional factors, other than solely expression levels, that delineate whether a tumor is dependent on a particular kinase. Approaches can include the use of phosphoproteomic or gene expression profiling analysis, examples of the latter being used to identify a kinase transformation fingerprint in human colorectal cancer and to define different prognoses, and efficacy of treatment has already been reported [12–14].

Tumors are dependent on neovascularization, the development of new blood vessels, to maintain a sufficient supply of nutrients and oxygen, which has led to the design of inhibitors directed against kinases that regulate this process. The clinical success of bevacizumab, an antibody that inhibits vascular endothelial growth factor receptor (VEGFR) signaling, has provided proof of concept for this approach [15]; however, additional targets on endothelial cells and other components of the vascular architecture are also being investigated.

Selectivity & approaches

The successful completion of the human genome has led to the advent of the kinome, currently containing more than 520 members [16]. It is apparent that any successful inhibitor should specifically target the intended kinase without affecting closely related subfamily members. The preponderance of RTKs as cancer targets has facilitated the development of therapeutic antibodies that inhibit receptor activation, either by preventing ligand–receptor interaction or by binding the receptor itself, inducing receptor internalization and, in parallel, causing cell killing via an antibody-dependent cellular cytotoxicity

(ADCC) response. Genetic engineering has enabled the development of fully humanized antibodies that overcome potential anti-antibody reactions and increase clinical efficacy. In a further level of sophistication, bispecific antibodies that recognize two different antigens have been developed. For example, MDX-447 and MDX-H210 (Medarex) contain one domain that recognizes either EGFR or HER-2, respectively, and another that is specific for CD64, thereby directing cytotoxic effector cells to receptor-overexpressing target cells. MDX-447 has entered Phase II clinical trials for the treatment of head and neck cancers that overexpress EGFR [17–19]. Some of the antibodies currently in clinical development are presented in TABLE 1.

Clinical agents derived from naturally occurring compounds comprise those directed towards the ATP-binding site and those with alternative mechanisms of action. Examples include flavopiridol (FPD/L868275/HMR1275), a semisynthetic flavone that associates with the ATP-binding site of cyclin-dependent kinases (CDKs) [20], and derivatives of the macrolide antibiotic rapamycin (sirolimus/Rapamune[®]), CCI-779 and Rad001 (everolimus/Certican[®]) that complex with the 12-kDa immunophilin FK506-binding protein (FKBP)-12, thereby specifically inhibiting the target of rapamycin (TOR) serine/threonine kinase [21].

In contrast, rationally designed low-molecular-weight compounds currently in the clinic are almost exclusively directed against the ATP-binding site of the kinase. ATP interacts, predominantly, via lipophilic/van der Waals interactions, within a cleft formed between two lobes of the kinase. Although it was initially felt that the conserved nature of the kinase catalytic domain would render identification of specific inhibitors difficult, alternative strategies, including attempts to design compounds that abrogate receptor–ligand interaction or prevent phosphotyrosine–Src homology (SH)-2 domain binding, have not met with clinical success. Cocrystallization of kinases with ATP analogs or inhibitors, as well as homology modeling studies, have been used to identify key structural features of ATP-binding sites that permit the design of relatively specific inhibitors [22]. As it appears improbable that completely specific inhibitors will be developed (i.e., one inhibitor, one kinase), attention has been focused on obtaining therapeutically beneficial clinical use in the absence of limiting toxicities.

Another method to obtain specificity within the ATP-binding site has been to target the inactive conformation of a kinase. Crystallization studies of different kinases have suggested that, whereas the active 3D conformation may be constrained due to catalytic necessity, greater diversity is observed in the structure of the inactive conformation – a phenomenon that presumably also reflects the diverse mechanisms by which kinase activity needs to be biologically regulated [23]. PD-173955 is a more potent but less specific inhibitor of Abl than imatinib. Solving the crystal structure of Abl in complex with either inhibitor suggests that this is due to the fact that PD-173955 is able to interact with multiple (active and inactive) conformations of the kinase, whereas imatinib specifically freezes the kinase into

Table 1. Examples of antibodies currently in clinical development.

Agent	Company	Structure	Target
Cetuximab (IMC-225)	ImClone/Merck/ Bristol-Myers Squibb	Chimeric monoclonal	EGFR
Trastuzumab	Genentech	Humanized monoclonal	HER-2
Pertuzumab	Genentech	Humanized monoclonal	HER-2
MDX-447	Medarex/Merck	Humanized bivalent monoclonal	EGFR
MDX-H210	Medarex/Novartis	Humanized bivalent monoclonal	HER-2
Bevacizumab	Genentech	Humanized monoclonal	VEGF
IMC-1C11	ImClone	Chimeric monoclonal	VEGFR2

EGFR: Epidermal growth factor receptor; VEGF(R): Vacular endothelial growth factor (receptor).

an inactive conformation. As discussed in the following section, one consequence of the specificity gained by imatinib binding the inactive kinase conformation is the risk of emergence of point mutations that can convey resistance to the drug without compromising catalytic activity. It is predicted that resistance mutations would be less likely to arise with compounds directed against the active conformation, as these would also perturb the catalytically active 3D kinase structure [24]. A representative sample of kinase inhibitors currently in the clinic is provided in TABLE 2.

Clinical examples of kinase inhibitors

Imatinib

The Philadelphia chromosome, which results in fusion and constitutive activation of the intracellular kinase Abl with Bcr (Bcr-Abl), is present in 95% of patients with CML and 15–30% of patients with acute lymphoblastic leukemia (ALL). Bcr-Abl is the unique cytogenetic aberration in chronic phase CML and induces a CML-like disease when expressed in murine bone marrow [25]. Imatinib, a potent inhibitor of Abl, platelet-derived growth factor receptor (PDGFR) and c-Kit, blocked Bcr-Abl-mediated proliferation of established cell lines and freshly isolated CML and ALL blasts [26,27]. Clinical response of patients at all stages of CML resulted in the approval of imatinib by the US Food and Drug Administration (FDA) for treatment of this indication [27,28]. Subsequently, imatinib was also approved for the treatment of GIST, a rare, chemoresistant sarcoma [29,30]. Imatinib is clinically effective in GIST due to the expression of constitutively activated c-Kit or, more infrequently, PDGFR α mutants [31,32]. Subsequently, other malignancies that display constitutive activation of the PDGFR, either due to the presence of aberrantly expressed

ligand, the COL1A1-PDGFR fusion in dermatofibrosarcoma protuberans [33,34], or fusion of the receptor TEL-PDGFR in chronic myelomonocytic leukemia (CML) [35] and Fip1L1-PDGFR α in hypereosinophilic syndrome (HES), have responded to treatment with imatinib [36]. Imatinib is additionally being considered for treatment of other neoplasms characterized by the presence of c-Kit mutants, such as seminoma [37] and systemic mast cell disease [38].

Issues in the clinic: emergence of resistance

Notably, although a stable response has been observed in chronic phase CML patients, most patients in accelerated or blast crisis, after an initial response, developed resistance to imatinib and relapsed [39]. In a similar manner, initial response of HES and GIST patients has been followed by the emergence of imatinib-resistant disease [40,41]. It has been suggested that resistance is an inevitable consequence of the treatment of genetically unstable disease with a single molecularly targeted agent. Several nonexclusive mechanisms by which resistance could occur can be envisaged: target amplification; mutation of the target to prevent action of the inhibitor; activation of a complementary pathway that bypasses requirement for the target; and upregulation of mechanisms that lower the intracellular concentrations of the inhibitor. However, analysis of CML patients, who have relapsed on treatment with imatinib, indicated that at least half of relapsed patients display point mutations within the kinase domain of Bcr-Abl [42,43]. Experimental recapitulation of these mutations has demonstrated that, to a greater or lesser extent, they render kinase activity resistant to the effects of imatinib [44]. The detection of mutations in two untreated blast phase CML patients, who subsequently failed to respond to imatinib, suggests that these mutations were not generated by exposure to the drug. Instead, it appears likely that a population of CML blasts contains multiple, albeit rare, mutant clones that are expanded by virtue of the resistance of their catalytic activity to the presence of imatinib [45,46]. The isolation of Fip1L1-PDGFR containing a Thr674Ile mutation, analogous with Thr315Ile imatinib-resistant mutation observed in Bcr-Abl, from an HES patient who relapsed after treatment with drug, suggests that the appearance of resistance mutations is not a peculiarity of Bcr-Abl or CML patients [36].

As described previously, imatinib binds to an inactive conformation of Abl, fitting in a pocket normally occupied by the DFG activation loop in the catalytically active kinase conformation [47,48]. After the initial finding of the Thr315Ile mutation, reported by Gorre and coworkers, over 30 different mutations have been identified in imatinib-resistant patients. Mapping the drug-resistant point mutations suggests that whereas some affect residues that make direct contact with imatinib, others convey resistance by sterically hindering the ability of the kinase to achieve an inactive conformation [46].

Experimental mutagenesis of Bcr-Abl kinase was used to identify amino-acid substitutions that conveyed resistance to imatinib [49]. Over 90 different mutations were identified, including all of the

Table 2. Examples of small-molecule kinase inhibitors in clinical development.

Agent	Company	Target	Indication
Gefitinib	AstraZeneca	EGFR	NSCLC
Erlotinib	OSI/Genentech/Roche	EGFR	NSCLC
Valatanib (PTK787/ ZK222584)	Novartis/Schering AG	VEGFR	Solid tumors
SU11248	Pfizer	VEGFR c-Kit Flt-3	Solid tumors GIST AML
Imatinib	Novartis	ABL PDGFR c-Kit	CML CMML, HES GIST
BMS-354825	Bristol-Myers Squibb	Src, Abl	Solid tumors
Rapamycin/ CCI-779	Wyeth	TOR	Solid tumors
SU011248	SUGEN/Pfizer	PDGF-R, Flt3, c-Kit, VEGFR	GIST
RAD001	Novartis	TOR	Solid tumors
Midostaurin/ PKC412	Novartis	Flt3, c-Kit, PDGFR	AML Solid tumors
PD 0325901	Pfizer	MEK	Solid tumors
BAY 43-9006	Bayer/Onyx	BRAF, VEGFR1-3	Malignant melanoma Renal cell carcinoma

AML: Acute myeloid leukemia; CML: Chronic myeloid leukemia; CMML: Chronic myelomonocytic leukemia; EGFR: Epidermal growth factor receptor; GIST: Gastrointestinal stromal tumor; HES: Hypereosinophilic syndrome; NSCLC: Non-small cell lung cancer; PDGFR: Platelet-derived growth factor receptor; VEGFR: Vascular endothelial growth factor receptor.

patient-derived imatinib-resistance mutations. Interestingly, a large number of mutations were found outside the kinase domain. As most studies of patient-derived blasts have been restricted to this domain, it is presently unclear whether these mutations represent experimental artifacts or would be compatible with the biologic ability of Bcr-Abl to drive CML. These studies also indicate that it should be possible to predict which mutated clones will arise in response to treatment with a particular drug, although the large number of identified mutations implies that it will be very difficult to identify a single therapeutic agent that could overcome all cases of resistance. Currently, attention is being focused on design of inhibitors that can overcome individual or a series of resistance mutations. More potent imatinib-like inhibitors could be used against those mutants that affect the conformation adopted by the kinase as it binds to the drug or those that occur outside the kinase domain. Similarly, such inhibitors should be effective against

resistance due to amplification or overexpression of Bcr-Abl. These inhibitors are not expected to be active against mutants that alter residues that directly interact with imatinib, such as the Thr315Ile mutant. Instead, inhibitors that directly target the active conformation may be useful, and several examples of this are known: PD-180970 has been shown to inhibit imatinib-resistant mutants of Bcr-Abl [50], and the Thr674Ile imatinib-resistant mutant of FipIL1-PDGFR was shown to be more sensitive to the staurosporine analog PKC412 than wild type Fip1L1-PDGFR [51]. Recently, BMS-354825, a dual Src/Abl inhibitor which is a 100-fold more potent inhibitor of Abl than imatinib, has been demonstrated to inhibit many tumor-derived, imatinib-resistant Abl mutants in preclinical mouse studies (Thr315Ile being a notable exception) [52]. Although the lack of selectivity of such agents may restrict their use in the clinic, a promising clinical response has recently been reported upon treatment of imatinib-resistant GIST with SU011248, an orally active inhibitor that targets several kinases, including PDGFR, Flt-3, c-Kit and VEGFR. More recent publications have demonstrated that the molecular mechanisms underlying imatinib resistance in GIST are similar to those observed in CML [53], suggesting that SU011248 acts via inhibition of imatinib-resistant c-Kit and PDGFR point mutants. However, additional effects via its action on compensatory pathways that are deregulated in the resistant tumors cannot be discounted [54].

Targeting parallel or downstream signaling pathways may represent a general approach by which resistance may be overcome. Bcr-Abl transformation results in activation of mitogen-activated protein kinase (MAPK), Akt and c-Src pathways. Inhibitors for these pathways are either in preclinical or clinical development and there is already evidence of synergistic or additive effects in combination with imatinib in model systems [55-57]. Increased understanding of the factors that modulate the effectiveness of combinational therapy, together with early identification and characterization of emerging resistance, are imperative steps in the design of more effective therapy.

EGFR inhibitors

The EGFR family of receptors consists of four structurally related transmembrane RTKs [58-61]. Ligand binding induces receptor hetero- and homodimerization and activation of the intrinsic tyrosine kinase domain. ErbB2/HER-2, that lacks any known ligand, is the preferred partner for the other receptors, however, the dimerization partner is also ligand and cell type dependent. Kinase activation results in initiation of downstream signaling cascades such as the PI3K/protein kinase B and MAPK pathways [60]. Overexpression of EGFR family members is frequently observed in solid tumors [62], whereas receptor mutations are less frequent, with the notable exception of brain tumors where both amplification and activating extracellular domain mutations have been reported. Mutation or high expression levels of EGFR and HER-2 are correlated with poor prognosis and more aggressive disease, and has led to intense efforts in identifying and developing therapeutic entities directed against these kinases [63,64].

Amongst these agents, the most advanced include small-molecule inhibitors (e.g., gefitinib and erlotinib) and therapeutic antibodies (e.g., trastuzumab, pertuzumab and cetuximab). Trastuzumab, a humanized immunoglobulin G1 antibody directed against HER-2, was the first FDA-approved monoclonal antibody directed against solid tumors. In a pivotal Phase III trial of patients with HER-2-positive metastatic breast cancer, combination of trastuzumab with chemotherapy significantly increased response, median duration of response and overall survival when compared with chemotherapy alone [65,66]. Trastuzumab has also demonstrated activity as a first-line single-agent therapy of patients with HER-2/3-overexpressing metastatic breast carcinoma [10]. Adverse effects have been documented, including cardiac dysfunction and narrowing of blood vessels, particularly when trastuzumab has been used in combination with anthracyclins [67].

Trastuzumab and pertuzumab bind to different epitopes on the extracellular domain of HER-2 [68]. Although both antibodies induce ADCC, pertuzumab fails to inhibit shedding of the intracellular domain of the receptor [69]. Interestingly, unlike trastuzumab, pertuzumab is effective in inhibiting HER-2 signaling in cell lines that do not overexpress the receptor [70]. This effect does not appear to be due to ADCC effects as the pertuzumab Fab domain is an equally effective inhibitor. Instead, pertuzumab binding blocks HER-2 signaling by preventing heterodimerization with coreceptors [68]. As expected, pertuzumab is effective both in the context of HER-2 overexpression, as well as in those tumors where HER-2 is transactivated via ligand-mediated activation of coreceptors (e.g., HER-3 and EGFR) [71,72]. Pertuzumab has shown an acceptable safety profile and has progressed to Phase II clinical trials [68].

Cetuximab is a humanized antibody that competes with ligand to bind the EGFR [73]. It has been reported that antibody binding results in receptor internalization without kinase activation. The consequent downregulation of signaling is associated with increased apoptosis, however, enhanced ADCC has also been reported [74,75]. The results from a Phase II clinical study led to an approval by the FDA for the use of cetuximab in combination with irinotecan in the treatment of patients with EGFR-expressing, metastatic colorectal cancer who are refractory or intolerant to chemotherapy. Cetuximab has also shown activity in patients with advanced pancreatic cancer upon combination with gemcitabine, a deoxycytidine analog, that represents the standard first-line chemotherapy for pancreatic cancer. Adverse effects observed upon administration of cetuximab included infusion symptoms as well as frequent (75%) occurrence of an acne-like rash, a common reaction to treatment with EGFR inhibitors. Rash could be used as a predictor of the survival of patients, the patients with the most intense rash deriving the greatest benefit from treatment [76].

Both gefitinib (EGFR inhibitor) and erlotinib (EGFR/HER-2 inhibitor) displayed activity against a wide spectrum of cell lines, as well as inhibiting growth and angiogenesis of tumor xenografts [77–80]. Activity hints were observed in Phase I clinical trials in patients with tumors

expected to express high EGFR levels, including colorectal cancer, non-small cell lung cancer (NSCLC) and head and neck squamous cell carcinoma (HNSCC) [81,83]. Similar objective response rates, approximately 10%, were observed in single-agent Phase II trials of both gefitinib and erlotinib in chemotherapy-refractory NSCLC patients. Although response rates were low, they were frequently durable and rapid in onset. This, in combination with an acceptable safety profile, led to the approval of gefitinib for treatment of NSCLC patients who had previously undergone chemotherapy, first in Japan, where a particularly high (19%) response rate was observed, and subsequently by the US and Australian regulatory authorities [84]. Unfortunately, Phase III trials in which gefitinib or erlotinib was combined with chemotherapy failed to demonstrate any improvement in response rate, progression-free or overall survival in NSCLC patients when compared with treatment with chemotherapy alone [80,85].

Issues in the clinic: patient selection

A large number of studies have been carried out to identify the reasons underlying the low patient response in trials with EGFR antagonistic agents. Post-treatment analysis of patient samples failed to demonstrate a clear correlation between the expression of EGFR and response to these drugs, thereby underlining a central problem in design of clinical trials – a lack of clear definition of which patients would benefit from kinase inhibitor therapy. In a similar manner, although a clear correlation existed between expression levels of HER-2 and response to trastuzumab monotherapy, less than a third of patients with receptor-positive malignancy benefited from treatment. The advent of validated antibodies that specifically recognize phosphorylated, and therefore activated, receptors has provided a potential explanation for this observation. Wide, tumor type-dependent variations in the percentage of tumors that contained phosphorylated receptor were observed in a post-treatment study of HER-2-overexpressing breast carcinomas [86,87]. Progression-free survival of patients with tumors containing phosphorylated HER-2 was found to be significantly longer after treatment with trastuzumab, when compared with patients with tumors lacking phosphoreceptor. Although these studies highlight the potential use of such an approach to identify tumors containing an activated kinase, it is apparent that intact transmission of the receptor signal to intracellular targets is also required. Resistance to gefitinib correlated with a failure to inhibit MAPK and Akt activation, despite blocking EGFR phosphorylation [88]. Further studies indicated that reconstitution of the tumor suppressor PTEN, a negative regulator of the PI3K/Akt pathway, restored sensitivity to EGFR inhibition [89]. Greater characterization of the activation state of an overexpressed kinase, its downstream signaling intermediates, as well as other signaling pathways that impinge on its action, are therefore imperative in defining a population of patients that would respond to an inhibitor. In the absence of a clear understanding of these factors, subpopulations of patients that do benefit from treatment risk being overlooked.

Importantly, two recent publications have provided the first instance of biomarkers that may be used to define tumor sensitivity to gefitinib [8,9]. Sequencing the EGFR tyrosine kinase domain identified a series of heterozygous point mutations that correlated with responsiveness of tumors to gefitinib. The presence of mutations mirrored the documented sensitivity to gefitinib, being more common in Japan, in adenocarcinomas, in women and in nonsmoking patients. Mutated EGFR displayed prolonged ligand-stimulated activation and increased sensitivity to gefitinib.

Although these studies provide hope that it will be possible to define a population of patients likely to be responsive to treatment with gefitinib, they raise additional questions. Does the absence of EGFR mutations in a responsive patient mean that mutations of additional pathways also render cancers sensitive to the effects of the drug? Does the different responsiveness of Japanese and US patients reflect different etiology of the disease due to racial, or as has been suggested, cultural (dietary) differences? Examination of more samples, including those from a recently reported trial of erlotinib, is essential to strengthen the statistical significance of these studies and to provide some answers to these questions.

VEGFR

VEGF and its cognate receptors, VEGFRs, play an essential role in angiogenesis, the process by which capillaries sprout from established blood vessels [90]. Simple diffusion of nutrients and oxygen becomes insufficient as tumors grow beyond a certain size, necessitating the *de novo* establishment of a blood supply [91,92]. VEGF expression is elevated in hypoxic and hypoglycemic conditions, upon expression of oncogenes, such as Ras, or due to inactivation of tumor suppressor genes, such as p53 or the von Hippel-Lindau gene [93]. Elevated levels of VEGF have been associated with poorer prognosis and increased chance of metastasis in a number of tumors [93]. In preclinical models, blockage of VEGF signaling with inhibitory antibodies inhibited the growth of tumor xenografts and reduced the number of metastases [94]. The proven role for VEGF in the regulation of vessel permeability suggests that its inhibition would also decrease the elevated interstitial pressure observed within tumors. Decreased interstitial tumor pressure would enhance delivery and therefore the effectiveness of coadministered chemotherapeutic agents, as has been observed in xenograft models [95].

A number of approaches have been adopted in targeting VEGF signaling. These include the use of therapeutic antibodies such as bevacizumab, monoclonal antibodies such as IMC-1C11 [96], small-molecular-weight inhibitors such as SU5416, SU-6668, SU11248 [97–99] and vatalanib (PTK787/ZK222584) [100], as well as angiozyme, a stabilized ribozyme that targets the pre-messenger RNA of VEGFR1 [101]. The most clinically advanced of these is bevacizumab, which binds to VEGF, inhibiting interaction with its cognate receptors. In a Phase II trial on patients with metastatic clear cell kidney carcinoma, no alteration in primary tumor growth or overall patient survival was observed,

however, a significant difference in time to progression was observed in those patients receiving bevacizumab compared with placebo [102,103]. However, most trials have focused on the combination of bevacizumab with chemotherapeutic agents. Combination with capecitabine, a fluoropyrimidine carbamate that inhibits thymidylate synthase, or vinorelbine, a vinca alkaloid, resulted in increased objective response rates in patients with metastatic breast cancer. Increased response rates and time to disease progression were also observed when NSCLC patients were treated with a combination of high-dose bevacizumab and carboplatin/paclitaxel [15]. Bevacizumab has recently been approved by the FDA as a first-line treatment, in combination with 5-fluorouracil/leucovorin/irinotecan therapy, for treatment of patients with metastatic colorectal cancer following results of Phase III trials where combination of bevacizumab with chemotherapy significantly increased overall survival, objective response rate and progression-free survival [104–106]. In these studies, no surrogate markers of activity were reported, and it therefore remains to be clarified whether the effects of bevacizumab are due to direct inhibition of neovascularization or modulation of vascular permeability.

Issues in the clinic: surrogate end points

Traditionally, anticancer drugs have been clinically evaluated in three phases. Phase I trials focus on drug safety and identification of an optimal dose and schedule for subsequent trials. Successive small patient groups are given incrementally higher concentrations of drug until a maximum tolerated dose (MTD) can be identified. Phase II trials are designed to evaluate the effectiveness of the MTD of the drug by measuring clinical responses (i.e., decrease in tumor size following treatment). The drug then enters Phase III trials where its efficacy, in terms of patient survival, is compared with that of the currently accepted standard treatment for that tumor type.

In contrast to cytotoxic agents, kinase inhibitors may have little/no toxicity and stable disease, rather than reduction in tumor size, should be considered a successful therapeutic outcome. Although most kinase inhibitors have, to date, been assessed using a classic trial design, these peculiarities make renewal and adaptation of clinical trials methodology essential. To reflect the proposed mechanism of action of the agent being studied, it is more appropriate to consider the use of biologic end points and surrogate markers. Instead of a MTD, Phase I studies should be designed to define the dose of a drug that provides maximal or sufficient target inhibition, that is the maximal biologically effective dose (MBD). Currently, however, the technology required to assess a MBD for most kinase inhibitors in clinical tissue is lacking. Therefore, most kinase inhibitor programs are accompanied by a parallel effort in translational research to develop robust, predictive and well-controlled biomarker assays. Approaches include generation and validation of phosphorylation-specific antibodies to detect modification of kinase or substrate in clinical tissue by immunochemistry, as well as the use of proteomics and expression profiling to identify treatment/response-dependent signatures.

Imaging studies to assess changes in tumor metabolism or vascular architecture have also been developed as alternative methods to assay compound activity. Dynamic, contrast-enhanced molecular resonance imaging (DCE-MRI) studies have been utilized to demonstrate that treatment with anti-VEGF agents (bevacizumab and PTK787) induces alterations in vascular structure and permeability [107,108]. Analysis of ^{18}F -fluorodeoxyglucose uptake by positron emission tomography following treatment of GIST patients with imatinib correlates with the activity of this drug in this highly metabolically active tumor. Relapse in patients undergoing treatment with the drug is accompanied by the reappearance of areas of metabolically active tumor [109].

During Phase I trials with imatinib, a clear correlation between clinical efficacy and inhibition of Bcr-Abl kinase activity in circulating leukemic cells was observed. However, as many kinase inhibitors are directed against solid tumors, from which it is inconvenient and frequently impossible to obtain biopsies to directly assess target inhibition, surrogate markers of drug efficacy are of particular importance. Although serendipitous clinical observations, such as rash (EGFR inhibitors) or alteration in hair color (c-Kit inhibitors), are of use in predicting treatment outcome and confirming patient compliance, assays measuring target kinase activity in a surrogate tissue are being developed for most kinase inhibitors entering trials. Surrogate biomarkers can also be used to provide information about the magnitude and durability of response to an inhibitor (i.e., how much kinase activity needs to be inhibited for how long to be clinically effective). CI-1040/PD 184352 is a non-ATP competitive MEK inhibitor that displayed activity against a range of tumor xenografts [110]. The phosphorylation of a MEK target, MAPK, in patient blood and tumor specimens, was used as a biomarker to demonstrate an acceptable safety profile at doses sufficient to inhibit the kinase. A failure to maintain target inhibition was probably responsible for the reported lack of activity of CI-1040 as a single agent in Phase II trials. Subsequently, PD 0325901, a more potent and soluble second-generation analog of CI-1040, has been introduced into Phase I clinical trials.

Clearly, the use of surrogate markers requires sufficient data on how reproducible, as well as how representative, they are in defining the threshold of the biologic response in the tumor tissue. The rapamycin derivatives, CCI-779 and RAD001, are currently in clinical trials with patients with advanced solid cancers. The target of both drugs, TOR, regulates translation of specific mRNA species with central roles in control of cell size and proliferation. Both drugs display antiproliferative activity as single agents in a range of preclinical cellular and xenograft systems [21]. CCI-779, the clinically more advanced compound, was evaluated using a traditional dose-escalation approach in Phase I trials. Mild-to-severe toxicities, including evidence of hepatic impairment, vomiting and thrombocytopenia, were observed in heavily pretreated patients. Notably, in a second study, based on a weekly rather

than biweekly schedule, no clear correlation between dose and adverse events was observed [111]. Partial response was observed in four patients with diverse tumor types, including NSCLC and renal cell carcinoma. In a Phase II trial with renal cell carcinoma patients, based on these results, objective response in 7% of patients and stable disease in 40% has been reported [112]. In contrast, during preclinical studies with Rad001, a relationship between unbound compound concentration and inhibition of a TOR substrate, p70S6K1, in peripheral blood mononuclear cells (PBMCs) and implanted tumor was established [113]. In Phase I trials, groups of patients received a single concentration of Rad001, calculated to achieve plasma concentrations sufficient to inhibit TOR [114,115]. Assessment of p70S6K1 activity in patient PBMCs was used as a marker of compound activity. In corroboration of these results, inhibition of p70S6K1 activity in PBMCs was found to significantly correlate with time to progression in post-treatment studies with CCI-779 [116]. Although further studies that correlate kinase inhibition in patient-derived tumor and PBMCs are required, these reports provide an example of how assessment of a surrogate marker may be utilized to provide a relevant pharmacodynamic readout of the efficacy of a kinase inhibitor.

Information provided by biomarkers and surrogate markers should also be incorporated into the design of Phase II clinical trials. Previously, large sample sizes were required due to the necessity to treat randomized patient populations with histologically defined tumors with a range of drug doses to assess both activity and toxicity issues. In contrast, once the MBD of a kinase inhibitor has been established, these trials should be less cumbersome, treating smaller numbers of patients with cancers known/likely to be dependent on the kinase targeted with doses close to the MBD. The end points of these trials should also be modified; assessment of shrinkage in tumor size is no longer appropriate to agents that will be largely cytostatic. Instead, if the new drug is active, it will modify the expected natural history of the disease. This can be measured, within each patient, by comparing time to progression on the new drug with time to progression registered with the previous first-line treatment. However, the heterogeneity in growth of many solid cancers will still necessitate recruitment of sufficient numbers of patients to distinguish between those tumors responding to the drug and those that naturally grow slowly. As a further refinement, Ratain and coworkers have proposed the randomized discontinuation design [117]. In this trial design, all patients are initially treated with the agent then patients with stable disease are randomized into placebo and continuing therapy groups. By initially selecting a more homogenous population, fewer patients should be required to demonstrate the activity of a drug. BAY 43-9006 (Sorafenib), a Raf inhibitor ($\text{IC}_{50} < 10 \text{ nM}$) that also displays activity against VEGFR1-3, c-Kit and PDGR ($\text{IC}_{50} 10\text{--}160 \text{ nM}$), has shown activity in Phase I studies of a cohort of 163 patients with a range of solid tumors [118]. Seven patients with renal cell carcinoma were treated with BAY 43-9006. Out of

these patients, one displayed a partial response and five reached stable disease. Although the target kinase in these tumors remains unresolved, renal cell carcinoma rarely contains mutations in BRAF, a Phase II trial incorporating the randomized discontinuation design was initiated. Preliminary data published from the trial indicate that after treatment with BAY 43-9006, 30% of patients had stable disease, while 40% responded (defined as >25% reduction in tumor volume as assessed by computed tomography or magnetic resonance imaging) [119]. Patients with stable disease were then randomized according to the trial design. Those patients that progressed while receiving placebo were rechallenged with the drug, whereas those progressing with the drug were taken off study. Although the final results of this trial are awaited, the preliminary findings using this trial design have prompted interest in the use of BAY 43-9006 for the treatment of renal cell carcinoma, and have led to a further Phase III study which is due to complete accrual in 2005.

Summary & conclusions

Advances have been made both in identifying the role of kinases in malignancies and characterizing inhibitors that can block their activity in a therapeutically relevant manner. The clinical efficacy of the compounds described herein demonstrates the utility and potential of kinase inhibitors, but also delineates a series of ground rules necessary for success. Central to these is a thorough molecular characterization of the role a kinase plays in the pathophysiology of a particular cancer. This is obviously a great challenge and can only be achieved by analysis of statistically relevant numbers of clinically defined samples. Experience with imatinib, and the recently reported studies of tumors responsive to gefitinib, suggest that activating gain-of-function mutations are more predictive than over-expression in defining a fundamental role for a kinase in cancer development. However, it is also apparent that the cellular context plays an important role, as exemplified by the higher probability of relapse to imatinib treatment in blast crisis patients with ALL rather than CML [120].

The emergence of resistance may be an inevitable consequence of any clinically effective kinase inhibitor. Specific point mutations that render kinase activity insensitive to the presence of inhibitor may be more likely to occur upon treatment with conformation-specific drugs, such as imatinib. Biochemical and structural studies have elucidated how these mutations overcome the action of the inhibitor and suggest methods of overcoming resistance. One approach, namely the treatment with a less specific inhibitor that interacts with all conformations of the kinase, has been attempted with some success in GIST patients who relapsed on treatment with imatinib. However, it should be noted that the inherent genetic instability of advanced stage tumors is likely to result in the appearance of other mechanisms of resistance to these agents. Alternative approaches such as targeting kinases expressed on nontransformed cells such as VEGFR, Tie-2 or PDGFR on endothelial cells or fibroblasts, where the emergence

of resistance would be less likely, or targeting multiple mutant kinases by inhibiting the action of molecular chaperones such as heat shock protein 90, may provide clinical benefit in these circumstances.

Expert opinion

Personalized medicine & targeted therapy

The past 5 years of clinical experience with kinase inhibitors has underscored the problems associated with employing targeted therapy against a molecularly heterogeneous disease. Differences between histologically similar cancer types has necessitated accrual of larger patient populations and led to the risk of clinical trials becoming underpowered, since beneficial effects on small percentages of patients are masked by the lack of response in the majority of patients. Furthermore, heterogeneity within a single tumor can result in outgrowth of subclones that do not possess the altered kinase activity, resulting in only partial response to treatment with an inhibitor. Molecular analysis of tumor tissues, taken before and after treatment, is beginning to permit characterization of those factors that identify drug sensitivity. Inevitably, these studies will also result in subfractionation of patient populations reducing the numbers that would benefit from administration of a particular kinase inhibitor. On the other hand, these studies should also help to identify aberrant activation of other pathways that represent potential targets. The low toxicity inherent in the premise of targeted therapy should permit the combination of inhibitors of different pathways allowing the development of treatment protocols tailored for individual patients. Targeting multiple aspects of the pathophysiology of a disease has already met with success in HIV therapy and examples of the beneficial effects of combinations of kinase inhibitors have already been reported in preclinical studies [121]. Personalized treatment strategies should provide the maximum benefit for the patient in terms of progression-free survival and improvement in quality of life, but will inevitably be associated with huge increases in the cost of patient healthcare.

Five-year view

If the last 10 years has seen the validation of kinase inhibitors as weapons in the war on cancer, it can be anticipated that the next 5 years will see the rationalization of their use in those situations where they will be predicted to have the maximal beneficial effect. Development of robust and validated diagnostic techniques to detect kinase activation in biologically relevant tissue, together with more appropriate design of clinical trials, will allow more effective evaluation of the clinical activity of new molecular entities. Increases in the armament of validated kinase inhibitors will be accompanied by design of trials utilizing the combination of agents, based on a clear mechanistic understanding of their actions in an appropriate cancer tissue. The increased cost of such an approach would be predicted to be accompanied by significant improvement in the clinical effectiveness of cancer treatment.

Key issues

- Sufficiently specific inhibitors can be developed with an acceptable safety profile.
- The development of preclinical models/assays that accurately reflect the role of kinases in tumor development is needed.
- Appropriate patient selection is essential to identify patients who would receive maximum benefit from treatment with a kinase inhibitor.
- Robust, accurate and sensitive biomarkers/surrogate markers need to be incorporated into clinical trials.
- Heterogeneity of the tumor microenvironment will require patient-specific drug combinations as well as continual monitoring of the tumor in terms of development of resistance and outgrowth of different clones.

References

Papers of special note have been highlighted as:

• of interest

•• of considerable interest

- Blume-Jensen P, Hunter T. Oncogenic kinase signaling. *Nature* 411, 355–365 (2001).
- Hunt JL. Molecular mutations in thyroid carcinogenesis. *Am. J. Clin. Pathol.* 118(Suppl.), S116–S127 (2002).
- Kitamura Y, Hirota S, Nishida T. Gastrointestinal stromal tumors (GIST): a model for molecule-based diagnosis and treatment of solid tumors. *Cancer Sci.* 94, 315–320 (2003).
- Garcia PI, Adams GP, Sundareshan P *et al.* Expression of mutated epidermal growth factor receptor by non-small cell lung carcinomas. *Cancer Res.* 53, 3217–3220 (1993).
- Davies H, Bignell GR, Cox C *et al.* Mutations of the BRAF gene in human cancer. *Nature* 417, 949–954 (2002).
- Identification of high frequency of BRAF mutations in various cancers, especially melanoma.**
- Bardelli A, Parsons DW, Silliman N *et al.* Mutational analysis of the tyrosine kinome in colorectal cancers. *Science* 300, 949 (2003).
- Samuels Y, Wang Z, Bardelli A *et al.* High frequency of mutations of the PIK3CA gene in human cancers. *Science* 304, 554 (2004).
- Paez JG, Janne PA, Lee JC *et al.* EGFR mutations in lung cancer: correlation with clinical response to gefitinib therapy. *Science* 304, 1497–1500 (2004).
- Identification of previously uncharacterized epidermal growth factor receptor (EGFR) point mutations in patients sensitive to EGFR inhibitors.**
- Lynch TJ, Bell DW, Sordella R *et al.* Activating mutations in the epidermal growth factor receptor underlying responsiveness of non-small-cell lung cancer to gefitinib. *N. Engl. J. Med.* 350, 2129–2139 (2004).
- Identification of previously uncharacterized EGFR point mutations in patients sensitive to EGFR inhibitors.**
- Vogel CL, Cobleigh MA, Tripathy D *et al.* First-line Herceptin monotherapy in metastatic breast cancer. *Oncology* 61(Suppl. 2), 37–42 (2001).
- Rubin BP, Schuetze SM, Eary JF *et al.* Molecular targeting of platelet-derived growth factor B by imatinib mesylate in a patient with metastatic dermatofibrosarcoma protuberans. *J. Clin. Oncol.* 20, 3586–3591 (2002).
- Singh D, Febbo PG, Ross K *et al.* Gene expression correlates of clinical prostate cancer behavior. *Cancer Cell* 1, 203–209 (2002).
- Glinksy GV, Glinkii AB, Stephenson AJ *et al.* Gene expression profiling predicts clinical outcome of prostate cancer. *J. Clin. Oncol.* 113, 913–923 (2004).
- Malek RL, Irby RB, Guo QM *et al.* Identification of Src transformation fingerprint in human colon cancer. *Oncogene* 21, 7256–7265 (2002).
- Zondor SD, Medina PJ. Bevacizumab: an angiogenesis inhibitor with efficacy in colorectal and other malignancies. *Ann. Pharmacother.* 38, 1258–1264 (2004).
- Manning G, Whyte DB, Martinez R *et al.* The protein kinase complement of the human genome. *Science* 298, 1912–1934 (2002).
- Repp R, van Ojik HH, Valerius T *et al.* Phase I clinical trial of the bispecific antibody MDX-H210 (anti-FcγRI × antiHER-2/neu) in combination with Filgrastim (G-CSF) for treatment of advanced breast cancer. *Br. J. Cancer* 89, 2234–2243 (2003).
- Sridhar SS, Seymour L, Shepherd FA. Inhibitors of epidermal-growth-factor receptors: a review of clinical research with a focus on non-small-cell lung cancer. *Lancet Oncol.* 4, 397–406 (2003).
- Wallace PK, Romet-Lemonne JL, Chokri M *et al.* Production of macrophage-activated killer cells for targeting of glioblastoma cells with bispecific antibody to FcγRI and the epidermal growth factor receptor. *Cancer Immunol. Immunother.* 49, 493–503 (2000).
- Fischer PM, Gianella-Borradori A. CDK inhibitors in clinical development for the treatment of cancer. *Expert Opin. Investig. Drugs* 12, 955–970 (2003).
- Bjornsti MA, Houghton PJ. The tor pathway: a target for cancer therapy. *Nature Rev. Cancer* 4, 335–348 (2004).
- Noble MEM, Endicott JA, Johnson LN. Protein kinase inhibitors: insights into drug design from structure. *Science* 303, 1800–1805 (2004).
- Huse M, Kuriyan J. The conformational plasticity of protein kinases. *Cell* 109, 275–282 (2002).
- Cowan-Jacob SW, Guez V, Fendrich G *et al.* Imatinib (STI571) resistance in chronic myelogenous leukemia: molecular basis of the underlying mechanisms and potential strategies for treatment. *Mini Rev. Med. Chem.* 4, 285–299 (2004).
- Daley GQ, Van Etten RA, Baltimore D. Induction of chronic myelogenous leukemia in mice by the P210bcr/abl gene of the Philadelphia chromosome. *Science* 247, 824–830 (1990).
- Heinrich MC, Griffith DJ, Druker BJ *et al.* Inhibition of c-kit receptor tyrosine kinase activity by STI 571, a selective tyrosine kinase inhibitor. *Blood* 96, 925–932 (2000).
- Druker BJ, Sawyers CL, Kantarjian H *et al.* Activity of a specific inhibitor of the BCR-ABL tyrosine kinase in the blast crisis of chronic myeloid leukemia and acute lymphoblastic leukemia with the Philadelphia chromosome. *N. Engl. J. Med.* 344, 1038–1042 (2002).
- Savage DG, Antman KH. Imatinib mesylate: a new oral targeted therapy. *N. Engl. J. Med.* 346, 683–693 (2002).

- 29 Demetri GD, von Mehren M, Blanke CD *et al.* Efficacy and safety of imatinib mesylate in advanced gastrointestinal stromal tumors. *N. Engl. J. Med.* 347, 472–480 (2002).
- 30 van Oosterom AT, Judson I, Verweij J *et al.* Safety and efficacy of imatinib (STI571) in metastatic gastrointestinal stromal tumours: a Phase I study. *Lancet* 358, 1421–1423 (2001).
- 31 Heinrich MC, Corless CL, Demetri GD *et al.* Kinase mutations and imatinib response in patients with metastatic gastrointestinal stromal tumor. *J. Clin. Oncol.* 21, 4342–4349 (2003).
- 32 Heinrich MC, Corless CL, Duensing A *et al.* PDGFRA activating mutations in gastrointestinal stromal tumors. *Science* 299, 708–710 (2003).
- 33 Rubin BP, Schuetze SM, Eary JF *et al.* Molecular targeting of platelet-derived growth factor B by imatinib mesylate in a patient with metastatic dermatofibrosarcoma protuberans. *J. Clin. Oncol.* 20, 3586–3591 (2002).
- 34 Sawyers CL. Imatinib GIST keeps finding new indications: successful treatment of dermatofibrosarcoma protuberans by targeted inhibition of the platelet-derived growth factor receptor. *J. Clin. Oncol.* 20, 3568–3569 (2002).
- 35 Apperley JF, Gardembas M, Melo JV *et al.* Response to imatinib mesylate in patients with chronic myeloproliferative diseases with rearrangements of the platelet-derived growth factor receptor- β . *N. Engl. J. Med.* 347, 481–487 (2002).
- 36 Cools J, DeAngelo DJ, Gotlib J *et al.* A tyrosine kinase created by fusion of the PDGFRA and FIP1L1 genes as a therapeutic target of imatinib in idiopathic hypereosinophilic syndrome. *N. Engl. J. Med.* 348, 1201–1214 (2003).
- 37 Kemmer K, Corless CL, Fletcher JA *et al.* KIT mutations are common in testicular seminomas. *Am. J. Pathol.* 164, 305–313 (2004).
- 38 Pardanani A, Elliott M, Reeder T *et al.* Imatinib for systemic mast-cell disease. *Lancet* 362, 535–536 (2003).
- 39 Griffin JD. Resistance to targeted therapy in leukaemia. *Lancet* 359, 458–459 (2002).
- 40 Cools J, Stover EH, Wlodarska I *et al.* The FIP1L1-PDGFR α kinase in hypereosinophilic syndrome and chronic eosinophilic leukemia. *Curr. Opin. Hematol.* 11(1), 51–57 (2004).
- 41 Heinrich MC, Corless CL, Demetri GD *et al.* Kinase mutations and imatinib response in patients with metastatic gastrointestinal stromal tumor. *J. Clin. Oncol.* 21, 4342–4349 (2003).
- 42 Gorre ME, Mohammed M, Ellwood K *et al.* Clinical resistance to STI-571 cancer therapy caused by BCR-ABL gene mutation or amplification. *Science* 293, 876–880 (2001).
- **Identification of some of the underlying mechanism of imatinib resistance in chronic myeloid leukemia patients.**
- 43 Gorre ME, Sawyers CL. Molecular mechanisms of resistance to STI571 in chronic myeloid leukemia. *Curr. Opin. Hematol.* 9, 303–307 (2002).
- 44 Sawyers CL. Opportunities and challenges in the development of kinase inhibitor therapy for cancer. *Genes Dev.* 17, 2998–3010 (2003).
- 45 Shah NP, Nicoll JM, Nagar B *et al.* Multiple BCR-ABL kinase domain mutations confer polyclonal resistance to the tyrosine kinase inhibitor imatinib (STI571) in chronic phase and blast crisis chronic myeloid leukemia. *Cancer Cell* 2, 117–125 (2002).
- 46 Shah NP, Sawyers CL. Mechanisms of resistance to STI571 in Philadelphia chromosome-associated leukemias. *Oncogene* 22, 7389–7395 (2003).
- 47 Nagar B, Hantschel O, Young MA *et al.* Structural basis for the autoinhibition of c-Abl tyrosine kinase. *Cell* 112, 859–871 (2003).
- 48 Schindler T, Bornmann W, Pellicena P *et al.* Structural mechanism for STI-571 inhibition of Abelson tyrosine kinase. *Science* 289, 1938–1942 (2000).
- 49 Azam M, Latek RR, Daley GQ. Mechanisms of autoinhibition and STI-571/imatinib resistance revealed by mutagenesis of BCR-ABL. *Cell* 112, 831–843 (2003).
- 50 La Rosee P, Corbin AS, Stoffregen EP *et al.* Activity of the Bcr-Abl kinase inhibitor PD-180970 against clinically relevant Bcr-Abl isoforms that cause resistance to imatinib mesylate (Gleevec, STI571). *Cancer Res.* 62, 7149–7153 (2002).
- 51 Cools J, Stover EH, Boulton CL *et al.* PKC412 overcomes resistance to imatinib in a murine model of FIP1L1-PDGFR α -induced myeloproliferative disease. *Cancer Cell* 3, 459–469 (2003).
- 52 Shah NP, Tran C, Lee FY *et al.* Overriding imatinib resistance with a novel ABL kinase inhibitor. *Science* 305(5682), 399–401 (2004).
- 53 Chen LL, Trent JC, Wu EF *et al.* A missense mutation in KIT kinase domain 1 correlates with imatinib resistance in gastrointestinal stromal tumors. *Cancer Res.* 64(17), 5913–5919 (2004).
- 54 Desai J, Maki R, Heinrich MC *et al.* Activity and tolerability of the multi-targeted tyrosine kinase inhibitor SU011248 in patients (pts) with metastatic gastrointestinal stromal tumor (GIST) refractory to imatinib mesylate. *Proc. Am. Soc. Clin. Oncol. Gastrointest. Cancer Symp.* (2004) (Abstract 7).
- **This and [51,52] present evidence of the potential use of active conformation targeting kinase inhibitors to overcome imatinib resistance in the preclinical and clinical settings.**
- 55 Yu C, Krystal G, Varticovski L *et al.* Pharmacologic mitogen-activated protein/extracellular signal-regulated kinase/tyrosine kinase/mitogen-activated protein kinase inhibitors interact synergistically with STI571 to induce apoptosis in Bcr/Abl-expressing human leukemia cells. *Cancer Res.* 62, 188–199 (2002).
- 56 Neshat MS, Raitano AB, Wang HG *et al.* The survival function of the Bcr-Abl oncogene is mediated by Bad-dependent and -independent pathways: roles for phosphatidylinositol 3-kinase and Raf. *Mol. Cell. Biol.* 20, 1179–1186 (2000).
- 57 Warmuth M, Simon N, Mitina O *et al.* Dual-specific Src and Abl kinase inhibitors, PP1 and CGP76030, inhibit growth and survival of cells expressing imatinib mesylate-resistant Bcr-Abl kinases. *Blood* 101, 664–672 (2003).
- 58 Mendelsohn J, Baselga J. The EGF receptor family as targets for cancer therapy. *Oncogene* 19, 6550–6565 (2000).
- 59 Velu TJ. Structure, function and transforming potential of the epidermal growth factor receptor. *Mol. Cell. Endocrinol.* 70, 205–216 (1990).
- 60 Yarden Y, Sliwkowski MX. Untangling the erbB signalling network. *Nature Rev.* 2, 127–137 (2001).
- 61 Yarden Y. The EGFR family and its ligand in human cancer signalling mechanisms and therapeutic opportunities. *Eur. J. Cancer* 37, S3–S8 (2001).
- 62 Salomon DS, Brandt R, Ciardiello F *et al.* Epidermal growth factor-related peptides and their receptors in human malignancies. *Crit. Rev. Oncol. Hematol.* 19, 183–232 (1995).
- 63 Garcia PI, Adams GP, Sundareshan P *et al.* Expression of mutated epidermal growth factor receptor by non-small cell lung carcinomas. *Cancer Res.* 53, 3217–3220 (1993).

- 64 Libermann TA, Nusbaum HR, Razon N *et al.* Amplification, enhanced expression and possible rearrangement of EGF receptor gene in primary human brain tumours of glial origin. *Nature* 313, 144–147 (1985).
- 65 Hirsch FR, Langer CJ. The role of HER2/neu expression and trastuzumab in non-small cell lung cancer. *Semin. Oncol.* 31(1 Suppl. 1), 75–82 (2004).
- 66 Slamon DJ, Leyland-Jones B, Shak S *et al.* Use of chemotherapy plus a monoclonal antibody against HER2 for metastatic breast cancer that overexpresses HER2. *N. Engl. J. Med.* 344, 783–792 (2001).
- **Seminal paper documenting the activity of trastuzumab in combination with chemotherapy for the treatment of breast cancer.**
- 67 Perez EA, Rodeheffer R. Clinical cardiac tolerability of trastuzumab. *J. Clin. Oncol.* 22, 322–329 (2004).
- 68 Franklin MC, Carey KD, Vajdos FF *et al.* Insights into ErbB signaling from the structure of the ErbB2-pertuzumab complex. *Cancer Cell.* 5, 317–328 (2004).
- 69 Molina MA, Codony-Servat J, Albanell J *et al.* Trastuzumab (Herceptin), a humanized antiHER2 receptor monoclonal antibody, inhibits basal and activated HER2 ectodomain cleavage in breast cancer cells. *Cancer Res.* 61, 4744–4749 (2001).
- 70 Agus DB, Akita RW, Fox WD *et al.* Targeting ligand-activated ErbB2 signaling inhibits breast and prostate tumor growth. *Cancer Cell.* 2, 127–137 (2002).
- 71 Mendoza N, Phillips GL, Silva J *et al.* Inhibition of ligand-mediated HER2 activation in androgen-independent prostate cancer. *Cancer Res.* 62, 5485–5488 (2002).
- 72 Jackson JG, St Clair P, Sliwkowski MX *et al.* Blockade of epidermal growth factor or heregulin-dependent ErbB2 activation with the anti-ErbB2 monoclonal antibody 2C4 has divergent downstream signaling and growth effects. *Cancer Res.* 64, 2601–2609 (2004).
- 73 Baselga J, Pfister D, Cooper MR *et al.* Phase I studies of anti-epidermal growth factor receptor chimeric antibody C225 alone and in combination with cisplatin. *J. Clin. Oncol.* 18, 904–914 (2000).
- 74 Clynes RA, Towers TL, Presta LG *et al.* Inhibitory Fc receptors modulate *in vivo* cytotoxicity against tumor targets. *Nature Med.* 6, 443–446 (2000).
- 75 Humblet Y. Cetuximab: an IgG(1) monoclonal antibody for the treatment of epidermal growth factor receptor-expressing tumours. *Expert Opin. Pharmacother.* 5(7), 1621–1633 (2004).
- 76 Saltz LB, Meropol NJ, Loehrer PJ Sr *et al.* Phase II trial of cetuximab in patients with refractory colorectal cancer that expresses the epidermal growth factor receptor. *J. Clin. Oncol.* 22, 1201–1208 (2004).
- 77 Dancey J, Sausville EA. Issues and progress with protein kinase inhibitors for cancer treatment. *Nature Rev. Drug Discov.* 2, 296–313 (2003).
- 78 Dancey J. Epidermal growth factor receptor inhibitors in clinical development. *Int. J. Radiat. Oncol. Biol. Phys.* 58, 1003–1007 (2004).
- 79 Dancey JE, Freidlin B. Targeting epidermal growth factor receptor – are we missing the mark? *Lancet* 362, 62–64 (2003).
- 80 Herbst RS, Giaccone G, Schiller JH *et al.* Gefitinib in combination with paclitaxel and carboplatin in advanced non-small-cell lung cancer: a Phase III trial – INTACT 2. *J. Clin. Oncol.* 22, 785–794 (2004).
- 81 Rich JN, Reardon DA, Peery T *et al.* Phase II trial of gefitinib in recurrent glioblastoma. *J. Clin. Oncol.* 22, 133–142 (2004).
- 82 Sanders ML. Investigations into the mechanism for suramin as an inhibitor of cAMP-dependent protein kinase. *Proc. Am. Chem. Soc.* (1996) (Abstract BIOL-035).
- 83 Soulieres D, Senzer NN, Vokes EE *et al.* Multicenter Phase II study of erlotinib, an oral epidermal growth factor receptor tyrosine kinase inhibitor, in patients with recurrent or metastatic squamous cell cancer of the head and neck. *J. Clin. Oncol.* 22, 77–85 (2004).
- 84 Dancey J. Epidermal growth factor receptor inhibitors in clinical development. *Int. J. Radiat. Oncol. Biol. Phys.* 58, 1003–1007 (2004).
- 85 Genentech Press Release. Phase III trials of Tarceva(TM) plus chemotherapy in first-line non-small cell lung cancer do not meet primary efficacy end point. Genentech, CA, USA (2003).
- 86 DiGiovanna MP, Chu P, Davison TL *et al.* Active signaling by HER-2/neu in a subpopulation of HER-2/neu-overexpressing ductal carcinoma *in situ*. clinicopathological correlates. *Cancer Res.* 62, 6667–6673 (2002).
- 87 Thor AD, Liu S, Edgerton S *et al.* Activation (tyrosine phosphorylation) of ErbB-2 (HER-2/neu): a study of incidence and correlation with outcome in breast cancer. *J. Clin. Oncol.* 18, 3230–3239 (2000).
- 88 Janmaat ML, Kruyt FAE, Rodriguez JA *et al.* Response to epidermal growth factor receptor inhibitors in non-small cell lung cancer cells: limited antiproliferative effects and absence of apoptosis associated with persistent activity of extracellular signal-regulated kinase or Akt kinase pathways. *Clin. Cancer Res.* 9, 2316–2326 (2003).
- 89 She QB, Solit D, Basso A *et al.* Resistance to gefitinib in PTEN-null HER-overexpressing tumor cells can be overcome through restoration of PTEN function or pharmacologic modulation of constitutive phosphatidylinositol 3'-kinase/Akt pathway signaling. *Clin. Cancer Res.* 9, 4340–4346 (2003).
- 90 Ferrara N, Gerber HP, LeCouter J. The biology of VEGF and its receptors. *Nature Med.* 9, 669–676 (2003).
- 91 Bergers G, Benjamin LE. Tumorigenesis and the angiogenic switch. *Nature Rev. Cancer* 3, 401–410 (2003).
- 92 Folkman J. Anti-angiogenesis: new concept for therapy of solid tumors. *Ann. Surg.* 175, 409–419 (1972).
- 93 Rosen LS. Clinical experience with angiogenesis signaling inhibitors: focus on vascular endothelial growth factor (VEGF) blockers. *Cancer Control* 9, 36–44 (2002).
- 94 Warren RS, Yuan H, Matli MR *et al.* Regulation by vascular endothelial growth factor of human colon cancer tumorigenesis in a mouse model of experimental liver metastasis. *J. Clin. Oncol.* 95, 1789–1797 (1995).
- 95 Borgstrom P, Gold DP, Hillan KJ *et al.* Importance of VEGF for breast cancer angiogenesis *in vivo*: implications from intravital microscopy of combination treatments with an antiVEGF neutralizing monoclonal antibody and doxorubicin. *Anticancer Res.* 19, 4203–4214 (1999).
- 96 Posey JA, Ng TC, Yang B *et al.* A Phase I study of antikinase insert domain-containing receptor antibody, IMC-1C11, in patients with liver metastases from colorectal carcinoma. *Clin. Cancer Res.* 9, 1323–1332 (2003).
- 97 Fabbro D, Manley PW. Su-6668. SUGEN. *Curr. Opin. Investig. Drugs* 2, 1142–1148 (2001).
- 98 Mendel DB, Laird AD, Smolich BD *et al.* Development of SU5416, a selective small molecule inhibitor of VEGFR receptor tyrosine kinase activity, as an anti-angiogenesis agent. *Anticancer Drug Des.* 15, 29–41 (2000).

- 99 Mendel DB, Laird AD, Xin X *et al.* *In vivo* antitumor activity of SU11248, a novel tyrosine kinase inhibitor targeting vascular endothelial growth factor and platelet-derived growth factor receptors: determination of a pharmacokinetic/pharmacodynamic relationship. *Clin. Cancer Res.* 9, 327–337 (2003).
- 100 Wood JM, Bold G, Buchdunger E *et al.* PTK787/ZK 222584, a novel and potent inhibitor of vascular endothelial growth factor receptor tyrosine kinases, impairs vascular endothelial growth factor-induced responses and tumor growth after oral administration. *Cancer Res.* 60, 2178–2189 (2000).
- 101 Sandberg JA, Parker VP, Blanchard KS *et al.* Pharmacokinetics and tolerability of an anti-angiogenic ribozyme (ANGIOZYME) in healthy volunteers. *J. Clin. Pharmacol.* 40, 1462–1469 (2000).
- 102 Yang JC, Haworth L, Sherry RM *et al.* A randomized trial of bevacizumab, an antivascular endothelial growth factor antibody, for metastatic renal cancer. *N. Engl. J. Med.* 349, 427–434 (2003).
- 103 Kabbinnar F, Hurwitz HI, Fehrenbacher L *et al.* Phase II, randomized trial comparing bevacizumab plus fluorouracil (FU)/leucovorin (LV) with FU/LV alone in patients with metastatic colorectal cancer. *J. Clin. Oncol.* 21, 60–65 (2003).
- **First clinical demonstration of the activity of the angiogenesis inhibitor bevacizumab.**
- 104 Genentech Press Release. FDA approves Avastin, a targeted therapy for first-line metastatic colorectal cancer patients. Genentech, CA, USA (2004).
- 105 Willett CG, Boucher Y, di Tomaso E *et al.* Direct evidence that the VEGF-specific antibody bevacizumab has antivascular effects in human rectal cancer. *Nature Med.* 10, 145–147 (2004).
- 106 Hurwitz H, Fehrenbacher L, Novotny W *et al.* Bevacizumab plus irinotecan, fluorouracil, and leucovorin for metastatic colorectal cancer. *N. Engl. J. Med.* 350, 2335–2342 (2004).
- 107 Morgan B, Thomas AL, Drevs J *et al.* Dynamic contrast-enhanced magnetic resonance imaging as a biomarker for the pharmacological response of PTK787/ZK 222584, an inhibitor of the vascular endothelial growth factor receptor tyrosine kinases, in patients with advanced colorectal cancer and liver metastases: results from two phase I studies. *J. Clin. Oncol.* 21, 3955–3964 (2003).
- 108 Willett CG, Boucher Y, di Tomaso E *et al.* Direct evidence that the VEGF-specific antibody bevacizumab has antivascular effects in human rectal cancer. *Nature Med.* 10, 145–147 (2004).
- 109 Stroobants S, Goeminne J, Seegers M *et al.* ¹⁸F-FDG-positron emission tomography for the early prediction of response in advanced soft tissue sarcoma treated with imatinib mesylate (Gleevec). *Eur. J. Cancer* 39, 2012–2020 (2003).
- 110 Sebolt-Leopold JS, Dudley DT, Herrera R *et al.* Blockade of the MAPK pathway suppresses growth of colon tumors *in vivo*. *Nature Med.* 5, 810–816 (1999).
- 111 Raymond E, Alexandre J, Faivre S *et al.* Safety and pharmacokinetics of escalated doses of weekly intravenous infusion of CCI-779, a novel mTOR inhibitor, in patients with cancer. *J. Clin. Oncol.* 22, 2336–2347 (2004).
- 112 Atkins MB, Hidalgo M, Stadler WM *et al.* Randomized Phase II study of multiple dose levels of CCI-779, a novel mammalian target of rapamycin kinase inhibitor, in patients with advanced refractory renal cell carcinoma. *J. Clin. Oncol.* 22, 909–918 (2004).
- 113 Boulay A, Zumstein-Mecker S, Stephan C *et al.* Antitumor efficacy of intermittent treatment schedules with the rapamycin derivative RAD001 correlates with prolonged inactivation of ribosomal protein S6 kinase 1 in peripheral blood mononuclear cells. *Cancer Res.* 64, 252–261 (2004).
- 114 O'Donnell A, Faivre S, Judson I *et al.* A Phase I study of the oral mTOR inhibitor RAD001 as monotherapy to identify the optimal biologically effective dose using toxicity, pharmacokinetic (PK) and pharmacodynamic (PD) end points in patients with solid tumours. *Proc. Am. Soc. Clin. Oncol.* (2003) (Abstract 803).
- 115 Lane H, Tanaka C, Kovarik J *et al.* Preclinical and clinical pharmacokinetic/pharmacodynamic (PK/PD) modeling to help define an optimal biological dose for the oral mTOR inhibitor, RAD001, in oncology. *Proc. Am. Soc. Clin. Oncol.* (2003) (Abstract 951).
- 116 Peralba JM, deGraffenried L, Friedrichs W *et al.* Pharmacodynamic evaluation of CCI-779, an inhibitor of mTOR, in cancer patients. *Clin. Cancer Res.* 9, 2887–2892 (2003).
- 117 Rosner GL, Stadler W, Ratain MJ. Randomized discontinuation design: application to cytostatic antineoplastic agents. *J. Clin. Oncol.* 4478–4484 (2002).
- 118 Strumberg D, Vliotitis D, Moeller JG *et al.* Results of Phase I pharmacokinetic and pharmacodynamic studies of the Raf kinase inhibitor BAY 43-9006 in patients with solid tumors. *Int. J. Clin. Pharmacol. Ther.* 40(12), 580–581 (2002).
- 119 Ahmad T, Eisen T. Kinase inhibition with BAY 43-9006 in renal cell carcinoma. *Clin. Cancer Res.* 10(18), 6388S–6392S (2004).
- 120 Druker BJ, Sawyers CL, Kantarjian H *et al.* Activity of a specific inhibitor of the BCR-ABL tyrosine kinase in the blast crisis of chronic myeloid leukemia and acute lymphoblastic leukemia with the Philadelphia chromosome. *N. Engl. J. Med.* 344, 1038–1042 (1992).
- 121 De Clercq E. Antiviral drugs in current clinical use. *J. Clin. Virol.* 30(2), 115–133 (2004).

Affiliations

- Mark A Pearson, PhD
Laboratory Head, Oncology, Novartis Institutes for Biomedical Research Basel, Novartis Pharma AG, WKL.125.4.16, CH-4002, Basel, Switzerland
Tel.: +41 616 961 016
Fax: +41 616 963 835
mark_andrew.pearson@pharma.novartis.com
- Dorian Fabbro, PhD
Head, Signaling Pathways Oncology, Novartis Institutes for Biomedical Research, Basel, Novartis Pharma AG, WKL.125.4.10, CH-4002, Basel, Switzerland
Tel.: +41 616 967 681
Fax: +41 616 963 835
dorian.fabbro@pharma.novartis.com